



Compendium of Veterinary Products - US edition

MERCK ANIMAL HEALTH
Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Nobivac[®] Intra-Trac[®]₃

Intervet/Merck Animal Health

CANINE ADENOVIRUS TYPE 2-PARAINFLUENZA-BORDETELLA BRONCHISEPTICA VACCINE

Modified Live Virus and Avirulent Live Culture

FOR ANIMAL USE ONLY

**IN THE ABSENCE OF A VETERINARIAN-CLIENT-PATIENT RELATIONSHIP,
FEDERAL REGULATIONS PROHIBIT THE RELABELING, REPACKAGING,
RESALE OR REDISTRIBUTION OF THE INDIVIDUAL CONTENTS OF THIS
PACKAGE.**

THIMEROSAL FREE

DIRECTIONS FOR USE - PLEASE READ CAREFULLY

Nobivac[®] Intra-Trac[®]₃ vaccine is recommended for use as an aid in the prevention of disease associated with canine adenovirus type 2, canine parainfluenza virus and *Bordetella bronchiseptica* infection in healthy dogs 3 weeks of age or older. These agents have been implicated as playing a role in the etiology of the condition known as canine kennel cough.

Duration of Immunity (DOI) has been demonstrated to be at least one year for *Bordetella bronchiseptica*.

WHEN TO VACCINATE

Vaccinate dogs at 3 weeks of age or older. Annual revaccination with one dose is recommended. Following the advice of the American Veterinary Medical Association on Vaccination Principles, an appropriate revaccination program for individual animals should be made based upon veterinarian-client-patient relationships.


PREPARATION OF THE VACCINE

1. Rehydrate live vaccine to desired volume with the accompanying Sterile Diluent.*
2. Draw the vaccine back into the syringe.
3. Remove needle from syringe.
4. Apply nasal applicator tip to syringe.

DOSAGE AND ADMINISTRATION

1 Dose - *Standard dose is 1 mL administered intranasally. Vaccine may be rehydrated with less than 1 mL of diluent to minimize discomfort when vaccinating small dogs and puppies. For one 1 mL dose, rehydrate vaccine with 1 mL sterile diluent or for one 0.5 mL dose, rehydrate vaccine with 0.5 mL sterile diluent. Instill the rehydrated vaccine into one or both nostrils of dogs as illustrated below.

5 Dose - *Standard dose is 1 mL administered intranasally. Vaccine may be rehydrated with less than 5 mL of diluent to minimize discomfort when vaccinating small dogs and puppies. For five 1 mL doses, rehydrate vaccine with 5 mL sterile diluent or for five 0.5 mL doses, rehydrate vaccine with 2.5 mL sterile diluent. Instill the rehydrated vaccine into one or both nostrils of five dogs as illustrated below.



CAUTION

1. THIS PRODUCT IS DESIGNED FOR INTRANASAL USE ONLY WITH THE ENCLOSED APPLICATORS. Systemic reactions resulting from inadvertent intramuscular or subcutaneous injection have been reported. Symptoms may include vomiting, diarrhea, lethargy, inappetence, jaundice and death associated with liver failure. Localized tissue necrosis at the injection site has also been reported. If inadvertent injection occurs, monitor the dog closely. Supportive therapy including IV fluids and treatment with gentamicin, tetracycline, trimethoprim/sulfa, or amoxicillin/clavulanic acid may be indicated. If anaphylactoid reaction occurs, use epinephrine.

2. Post vaccinal reactions consisting of mild canine cough syndrome may occur following use of this vaccine.

3. For animal use only. Store at 2° - 7°C (35° - 45°F). Use entire contents when first opened. Burn this container and all unused contents.

4. This vaccine contains penicillin, streptomycin and nystatin as preservatives.

FOR INTRANASAL USE ONLY

For veterinary use in dogs only.

Intervet Inc., d/b/a Merck Animal Health, Omaha, NE 68103 USA

1 800 224-5318 (USA)

U.S. Veterinary License No. 165A

For patent information:

<http://www.merck.com/product/patent/home.html>

	Code
2x5 DOSE - VARIED DOSE VOLUME	065314 142176-02, 145868-02
25x1 DOSE - VARIED DOSE VOLUME	065313 163334-03
150x1 DOSE - VARIED DOSE VOLUME	065310 164837-03, 153303-03

CPN: 1047240.6

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 23, 2018
Copyright 2018 - Animalytix LLC